Table 8.1 ***Illicit Drug Use in Past Month and Illicit Drug Use Other Than Marijuana in Past Month*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Illicit Drug Use in Past Month		Illicit Drug Use Other Than Marijuana in Past Month	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	8.75	(7.55 - 10.12)	3.97	(3.15 - 5.00)
Kent	7.50	(5.77 - 9.69)	3.92	(2.79 - 5.48)
New Castle (excluding Wilmington City)	9.07	(7.56 - 10.85)	3.96	(3.00 - 5.22)
Sussex	7.61	(5.83 - 9.88)	3.75	(2.65 - 5.28)
Wilmington City	12.08	(8.99 - 16.05)	4.70	(3.12 - 7.03)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates are based on data from original questions, excluding those on the use of over-the-counter drugs or new methamphetamine items that were added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.2 ***Marijuana Use in Past Month, Average Annual Rate of First Use of Marijuana, and Perceptions of Great Risk of Smoking Marijuana Once a Month*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Marijuana Use in Past Month		Average Annual Rate of First Use of Marijuana[1]		Perceptions of Great Risk of Smoking Marijuana Once a Month	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	6.52	(5.58 - 7.62)	1.68	(1.46 - 1.94)	36.75	(34.07 - 39.52)
Kent	5.40	(4.03 - 7.20)	1.58	(1.25 - 2.01)	38.45	(33.71 - 43.42)
New Castle (excluding Wilmington City)	6.91	(5.69 - 8.36)	1.88	(1.56 - 2.26)	34.21	(30.84 - 37.74)
Sussex	5.60	(4.16 - 7.51)	1.16	(0.91 - 1.48)	41.37	(36.59 - 46.32)
Wilmington City	8.68	(6.30 - 11.85)	2.17	(1.59 - 2.95)	37.09	(31.34 - 43.23)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

[1] *Average annual marijuana initiation rate* = $100*\{[X_1 \div (0.5 * X_1 + X_2)] \div 2\}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of persons who never used marijuana. Both of the computation components, X_1 and X_2, are based on a survey-weighted hierarchical Bayes estimation approach. Note that the age group is based on a respondent's age at the time of the interview, not his or her age at first use. The prediction intervals for Average Annual Rate of First Use of Marijuana use a simultaneous solution for First Use of Marijuana and Never Used Marijuana outcomes, which had age group-specific random effects that were allowed to be correlated across the two outcomes. The associated Markov Chain Monte Carlo chains were used to calculate the posterior variance.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.3 ***Marijuana Use in Past Year, Cocaine Use in Past Year, and Nonmedical Use of Pain Relievers in Past Year*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Marijuana Use in Past Year		Cocaine Use in Past Year		Nonmedical Use of Pain Relievers in Past Year	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	11.12	(9.79 - 12.61)	2.56	(1.93 - 3.38)	5.00	(4.17 - 5.97)
Kent	8.91	(7.08 - 11.17)	2.27	(1.49 - 3.44)	5.14	(3.92 - 6.71)
New Castle (excluding Wilmington City)	11.47	(9.81 - 13.37)	2.66	(1.94 - 3.66)	5.16	(4.16 - 6.38)
Sussex	9.88	(7.75 - 12.51)	2.27	(1.49 - 3.42)	4.60	(3.45 - 6.12)
Wilmington City	16.37	(12.57 - 21.06)	3.17	(1.95 - 5.14)	4.72	(3.41 - 6.50)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.4 ***Alcohol Use in Past Month, Binge Alcohol Use in Past Month, and Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Alcohol Use in Past Month		Binge Alcohol Use in Past Month[1]		Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	54.85	(52.14 - 57.54)	23.56	(21.49 - 25.75)	40.92	(38.18 - 43.72)
Kent	51.27	(46.06 - 56.45)	23.04	(19.63 - 26.85)	42.10	(37.68 - 46.66)
New Castle (excluding Wilmington City)	59.00	(55.23 - 62.67)	24.72	(22.07 - 27.58)	39.46	(36.10 - 42.93)
Sussex	50.27	(45.23 - 55.31)	21.46	(18.23 - 25.08)	40.33	(36.12 - 44.68)
Wilmington City	48.67	(42.09 - 55.30)	22.82	(18.78 - 27.44)	48.57	(43.39 - 53.78)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

[1] Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.5 ***Alcohol Use in Past Month and Binge Alcohol Use in Past Month*** **in Delaware among Persons Aged 12 to 20, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Alcohol Use in Past Month among Persons Aged 12 to 20		Binge Alcohol Use in Past Month among Persons Aged 12 to 20[1]	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	29.49	(27.16 - 31.93)	20.08	(18.01 - 22.33)
Kent	29.13	(25.19 - 33.41)	18.20	(14.82 - 22.16)
New Castle (excluding Wilmington City)	32.22	(29.28 - 35.32)	22.86	(20.11 - 25.86)
Sussex	24.81	(21.05 - 28.98)	17.10	(13.72 - 21.11)
Wilmington City	22.73	(18.15 - 28.07)	12.87	(9.69 - 16.90)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

[1] Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.6 ***Cigarette Use in Past Month, Tobacco Product Use in Past Month, and Perceptions of Great Risk of Smoking One or More Packs of Cigarettes Per Day*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Cigarette Use in Past Month		Tobacco Product Use in Past Month[1]		Perceptions of Great Risk of Smoking One or More Packs of Cigarettes Per Day	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	25.31	(23.14 - 27.62)	29.10	(26.86 - 31.46)	74.68	(72.44 - 76.79)
Kent	26.43	(22.89 - 30.30)	30.28	(26.39 - 34.47)	72.92	(69.12 - 76.42)
New Castle (excluding Wilmington City)	23.09	(20.49 - 25.92)	26.49	(23.73 - 29.45)	76.75	(74.06 - 79.25)
Sussex	28.61	(24.65 - 32.93)	33.18	(29.25 - 37.35)	70.84	(66.86 - 74.53)
Wilmington City	28.13	(23.81 - 32.89)	32.20	(27.55 - 37.23)	75.23	(71.35 - 78.73)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

[1] Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.7 ***Alcohol Dependence in Past Year and Illicit Drug Dependence in Past Year*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Alcohol Dependence in Past Year		Illicit Drug Dependence in Past Year[1]	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	3.17	(2.57 - 3.92)	2.03	(1.59 - 2.60)
Kent	3.00	(2.18 - 4.12)	2.37	(1.59 - 3.51)
New Castle (excluding Wilmington City)	3.19	(2.43 - 4.17)	1.93	(1.39 - 2.67)
Sussex	3.02	(2.18 - 4.15)	1.87	(1.23 - 2.83)
Wilmington City	3.79	(2.66 - 5.38)	2.38	(1.46 - 3.87)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

[1] Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically; these estimates are based on data from original questions, excluding those on the use of over-the-counter drugs or new methamphetamine items that were added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.8 ***Alcohol Dependence or Abuse in Past Year, Illicit Drug Dependence or Abuse in Past Year, and Dependence on or Abuse of Illicit Drugs or Alcohol in Past Year*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Alcohol Dependence or Abuse in Past Year		Illicit Drug Dependence or Abuse in Past Year[1]		Dependence on or Abuse of Illicit Drugs or Alcohol in Past Year[1]	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	6.81	(5.83 - 7.94)	3.01	(2.43 - 3.73)	8.71	(7.61 - 9.95)
Kent	6.33	(4.97 - 8.04)	3.28	(2.32 - 4.62)	8.49	(6.77 - 10.59)
New Castle (excluding Wilmington City)	7.01	(5.78 - 8.48)	2.82	(2.12 - 3.74)	8.74	(7.31 - 10.40)
Sussex	6.29	(4.89 - 8.06)	3.11	(2.15 - 4.47)	8.18	(6.46 - 10.32)
Wilmington City	7.83	(5.96 - 10.21)	3.34	(2.15 - 5.15)	10.28	(7.88 - 13.31)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

[1] Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically; these estimates are based on data from original questions, excluding those on the use of over-the-counter drugs or new methamphetamine items that were added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 8.9 ***Needing But Not Receiving Treatment for Alcohol Use in Past Year and Needing But Not Receiving Treatment for Illicit Drug Use in Past Year*** **in Delaware among Persons Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**

State/Substate Region	Needing But Not Receiving Treatment for Alcohol Use in Past Year[1]		Needing But Not Receiving Treatment for Illicit Drug Use in Past Year[2]	
	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Delaware	6.40	(5.46 - 7.50)	2.80	(2.25 - 3.49)
Kent	5.95	(4.61 - 7.64)	3.24	(2.25 - 4.65)
New Castle (excluding Wilmington City)	6.70	(5.51 - 8.13)	2.46	(1.85 - 3.26)
Sussex	5.79	(4.43 - 7.54)	3.10	(2.09 - 4.59)
Wilmington City	7.03	(5.26 - 9.35)	3.23	(2.08 - 5.00)

NOTE: For Substate Region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health*.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach, and the 95 percent prediction (credible) intervals are generated by Markov Chain Monte Carlo techniques.

[1] Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but not receiving treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], or mental health centers).

[2] Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but not receiving treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], or mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, these estimate are based on data from original questions excluding those on the use of over-the-counter drugs or new methamphetamine items that were added in 2005 and 2006. See Section B.4.8 in the Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.1.1 ***Illicit Drug Use in Past Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.1.2 ***Illicit Drug Use Other Than Marijuana in Past Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.2.1 ***Marijuana Use in Past Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.2.2 ***First Use of Marijuana*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Average Annual Rates Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.2.3 ***Perceptions of Great Risk of Smoking Marijuana Once a Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.3.1 ***Marijuana Use in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.3.2 ***Cocaine Use in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.3.3 ***Nonmedical Use of Pain Relievers in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.4.1 ***Alcohol Use in Past Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.4.2 ***Binge Alcohol Use in Past Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.4.3 ***Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.5.1 ***Alcohol Use in Past Month among Persons Aged 12 to 20*** **in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.5.2 ***Binge Alcohol Use in Past Month among Persons Aged 12 to 20*** **in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.6.1 ***Cigarette Use in Past Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.6.2 ***Tobacco Product Use in Past Month*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.6.3 ***Perceptions of Great Risk of Smoking One or More Packs of Cigarettes Per Day*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.7.1 ***Alcohol Dependence in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.7.2 ***Illicit Drug Dependence in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.8.1 ***Alcohol Dependence or Abuse in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.8.2 ***Illicit Drug Dependence or Abuse in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.8.3 ***Dependence on or Abuse of Illicit Drugs or Alcohol in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.9.1 ***Needing But Not Receiving Treatment for Alcohol Use in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Figure 8.9.2 ***Needing But Not Receiving Treatment for Illicit Drug Use in Past Year*** **among Persons Aged 12 or Older in Delaware, by Substate Region: Percentages, Annual Averages Based on 2006, 2007, and 2008 NSDUHs**



NOTE: The legend's ranges were created by dividing 344 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see Section D of the report on *Substate Estimates from the 2006-2008 National Surveys on Drug Use and Health.*

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.